



Food, Energy, Water, Climate Nexus: Potential in Cameroon

Ackom, Emmanuel

Publication date:
2014

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Ackom, E. (2014). *Food, Energy, Water, Climate Nexus: Potential in Cameroon*. Paper presented at Nexus 2014: Water, Food, Climate and Energy Conference, Chapel Hill, NC, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Food, Energy, Water, Climate Nexus: Potential in Cameroon

Emmanuel Ackom, PhD

GNESD

UNEP Risø Centre

DTU Management Engineering

Nexus 2014 - Water, Food, Climate and Energy Conference
7th March, 2014, Chapel Hill, North Carolina.

Outline of today's presentation

Background on the GNESD Network (facilitated by UNEP)

Food, Energy, Water, Climate Nexus: Potential in Cameroon

- Food production
- Bioelectricity production potential from agricultural residues (20% use)
- Water savings (potential) relative to the use of crude oil electricity
- GHG emissions reduction potential relative to the use of crude oil electricity
- Concluding comments
- Acknowledging our donors/sponsors

What is GNESD?

GNESD:



launched at the World Summit on Sustainable Development
(2002)

is a global knowledge network involving 10 Centres of Excellence
and Network Partners.

Objectives of GNESD:

Knowledge network

Policy analysis on environmentally benign energy systems and services that:

- can help achieve Millennium Development Goals
- are not harmful to human health;
- do not conflict with our food supply;
- result in poverty alleviation and
- achieving sustainable development in member countries

Centres of Excellence from developing countries




- Energy Research Centre, Univ. of Cape Town, (South Africa)
- AFREPREN (Kenya)
- ENDA-TM(Senegal)
- Mediterranean Renewable Energy Centre MEDREC (Tunisia)
- Asian Institute of Technology (Thailand)
- TERI (India)
- Energy Research Institute (China)
- Fundación Bariloche (Argentina)
- CENBIO/Univ. of São Paulo & CENTROCLIMA/Fed. Univ. of Rio de Janeiro (Brazil)
- Molina Centre on Energy and Environment, Mexico

How GNESD works ...




- Network Centres cooperate through activity based working groups
- Multi-regional (or country) efforts and cross learning
- Annual assemblies, teleconferences etc
- A steering committee provides strategic direction and oversight
- Management structure
- UNEP affiliated secretariat based in Denmark



Selected Summary for Policy Makers (SPM) Publications: download (free) at www.gnesd.org



Selected SPM Publications: download (free) at www.gnesd.org



Publication (contd.):


Energy Policy 63 (2013) 101–113



Contents lists available at ScienceDirect

Energy Policy

journal homepage: www.elsevier.com/locate/enpol



Modern bioenergy from agricultural and forestry residues in Cameroon: Potential, challenges and the way forward



Emmanuel K. Ackom^{a,*}, Dieudonne Alemagi^{b,1}, Nana B. Ackom^{c,2}, Peter A. Minang^{d,3}, Zac Tchoundjeu^{e,4}

^a Global Network on Energy for Sustainable Development (GNESD), UNEP Risø Centre on Energy, Climate and Sustainable Development, Frederiksborgvej 399, Department of Management Engineering, Technical University Denmark, 4000 Roskilde, Denmark

^b World Agroforestry Centre Regional Office, PO Box 16317, Yaoundé, Cameroon

^c Faculty of Engineering Sciences, University of Ghana, PO Box LG25, Legon, Ghana

^d World Agroforestry Centre, PO Box 30677, Nairobi 00100, Kenya

^e World Agroforestry Centre Regional Office, PO Box 16317, Yaoundé, Cameroon

H I G H L I G H T S

- Environmentally benign residues amount to 1.11×10^6 bone dry tonnes per annum.
- 0.12–0.32 billion litres of bio ethanol annually to displace 18–48% national gasoline use.
- 0.08–0.22 billion litres of biomass to BTL diesel per year to offset 17–45% of diesel use.
- 0.76–2.02 TW h of electricity, representing 15–38% of Cameroon's consumption.
- Residues could offset only 3% of national consumption of traditional biomass.

Food, Energy, Water, Climate Nexus: Potential in Cameroon

- Food production
- Bioelectricity production potential from agricultural residues (20% use)
- Water savings (potential) relative to the use of crude oil electricity
- GHG emissions reduction potential relative to the use of crude oil electricity
- Concluding comments




Cameroon:



Map source: www.cigarinspector.com


- Total area of 475 440 km²
- 3 times the size of North Carolina
- 1/21 times size of USA

**Cameroon: Electricity Access = 48.7 % population
(in)accessibility = 51.3% population**



Source: IEA, 2009

Current Electricity Generation:



Total: 1016 MW

| ^a Agricultural crop residues | Production | Residue type | ^c Residue to product ratio (RPR) | Moisture content | ^e Lower heating Value | Residue | Residue | Residue, 20% sustainable extraction | Energy Potential (bone dry tons X MJ/kg) | ^h MW h, (GJ × 0.28 × efficiency) | |
|---|-----------------|--------------|---|------------------|----------------------------------|------------|-----------------|-------------------------------------|--|---|-----------------|
| Units | (tons) | n/a | n/a | (%) | (MJ/kg) | (wet tons) | (bone dry tons) | (bone dry tons) | GJ | 15% efficiency | 40% efficiency |
| | | | | | | | | | | MW h (Low) | MW h (High) |
| Maize | 1.67E+06 | Stalk | 1.5 | 15 | 15.48 | 2.51E+06 | 2.13E+06 | 4.27E+05 | 6.61E+06 | 2.78E+05 | 7.40E+05 |
| Sorghum | 9.00E+05 | Stalk | 2.62 | 15 | 17.00 | 2.36E+06 | 2.00E+06 | 4.01E+05 | 6.81E+06 | 2.86E+05 | 7.63E+05 |
| Rice | 1.75E+05 | Straw | 1.5 | 15 | 15.56 | 2.62E+05 | 2.23E+05 | 4.46E+04 | 6.94E+05 | 2.91E+04 | 7.77E+04 |
| Millet | 5.53E+04 | Stalk | 3 | 15 | 15.51 | 1.66E+05 | 1.41E+05 | 2.82E+04 | 4.37E+05 | 1.84E+04 | 4.90E+04 |
| Wheat | 9.00E+02 | Straw | 1.2 | 15 | ^f 15.60 | 1.08E+03 | 9.18E+02 | 1.84E+02 | 2.86E+03 | 1.20E+02 | 3.21E+02 |
| Sugarcane | 1.45E+06 | Bagasse | 0.3 | 75 | 13.38 | 4.35E+05 | 1.09E+05 | 2.18E+04 | 2.91E+05 | 1.22E+04 | 3.26E+04 |
| Cocoa | 2.64E+05 | Pods, Husk | 1 | 15 | 15.48 | 2.64E+05 | 2.24E+05 | 4.49E+04 | 6.95E+05 | 2.92E+04 | 7.78E+04 |
| Coconut | 5.00E+03 | Shell | 0.6 | 10 | 10.61 | 3.00E+03 | 2.70E+03 | 5.40E+02 | 5.73E+03 | 2.41E+02 | 6.42E+02 |
| Coffee | 6.66E+04 | Husk | 2.1 | 15 | 12.56 | 1.40E+05 | 1.19E+05 | 2.38E+04 | 2.99E+05 | 1.25E+04 | 3.34E+04 |
| Sub-total | 4.59E+06 | | | | | | | 9.92E+05 | | 6.66E+05 | 1.78E+06 |

Food

Energy

^a Agricultural crop production based on year 2010 statistics information (FAOSTAT, 2012).

^c Residue to product ratio (RPR) was based on published information (OECD/IEA, 2010), except for industrial roundwood RPR.

^e Lower heating values were based on published information (NREL, 2008), except for wheat, industrial roundwood and sawnwood.

^f Lower heating values on wheat were based on published information (Maas et al., 2008).

^h Decentralized bioelectricity generation method, based on Mendu et al., 2012.



Energy potential (bioelectricity) from residues:

- ***Best case:
33% of national electricity consumption***
- ***Least case:
13% of national electricity consumption***
- ***Residues could essentially power most farming communities at decentralized power system scales***




GHG emission reduction potential (bioelectricity) from residues - (reference to crude oil powered electricity)

➤ ***Best case:***
1.7 Mt CO₂

➤ ***Least case:***
0.6 Mt CO₂



Water: Estimated 2 to 8 billion litres/year potential savings



Source: www.ewb-dc.org

Conclusion

- *Bioelectricity from agricultural residues exhibit good food-energy-water-climate nexus*
- *Extending electricity access should not always be about long transmission lines but decentralized systems could play key roles especially in rural farming areas*
- *The knowledge could possibly help inform decision makers regarding the good potential of residues for social and environmentally benign development*

Acknowledgement – donor gov'ts/organizations

GNESD secretariat wish to thank:

- Government of Germany;
- Government of Denmark;
- Government of France;
- Government of Italy;
- Government of the United Kingdom;
- UN Foundation;
- UNDP (supported Energy Access activities & outreach activities in the RET theme)
- UNEP

THANK YOU

Emmanuel Ackom, Programme Manager,
UNEP Riso Centre
Email:emac@dtu.dk